

The Unusual Cataclysmic Binary Star RBS 0490 and the Space Density of Cataclysmics ¹

John R. Thorstensen

*Department of Physics and Astronomy
6127 Wilder Laboratory, Dartmouth College
Hanover, NH 03755-3528;
john.thorstensen@dartmouth.edu*

Sébastien Lépine and Michael Shara

*Department of Astrophysics, Division of Physical Science
American Museum of Natural History
Central Park West at 79th Street
New York, NY 10024*

ABSTRACT

RBS (Rosat Bright Source) 0490 is a cataclysmic variable star (CV) with unusually strong emission lines. The strength of the emission lines led to a suggestion that the object is intrinsically faint and correspondingly nearby (~ 33 pc), which, if true, would strongly affect estimates of the CV space density. Here we report astrometry, filter photometry, and time-series spectroscopy of this object. The astrometry gives an absolute parallax $\pi_{\text{abs}} = 4.5 \pm 1.5$ mas and a relative proper motion of 102 mas yr^{-1} . A Bayesian procedure gives a very uncertain distance estimate of $d \sim 300$ pc, and the small parallax alone implies $d > 133$ pc (at two standard deviations). The mean V magnitude is 17.4, which implies $M_V = 10.9 - 5 \log(d/200 \text{ pc})$, neglecting extinction. At 200 pc, the space velocity would be over 90 km s^{-1} with respect to the LSR. The time-series spectroscopy shows a possible emission-line radial-velocity period near 46 min. This would be unusually short for an orbital period and it may represent some other clock in the system.

Subject headings: stars – individual (RBS 490); binaries – close; novae, cataclysmic variables

1. Introduction

Cataclysmic variable stars (CVs) are close binaries in which a white dwarf accretes matter from a less-evolved companion (the secondary), which usually resembles a lower main sequence star. CVs have a rich phenomenology, and the theory of CVs ranges across wide areas of astrophysics; Warner (1995) presents a comprehensive review.

CVs are common enough that there are several with distance $d < 100$ pc, but their space

density remains somewhat controversial, in part because of distance uncertainties. Also, the completeness of the known sample is not perfectly understood. For example, all-sky camera surveys (e.g., Pojmański 2001) frequently turn up previously undiscovered, bright dwarf novae in outburst. Many new CVs have also been found in the Sloan Digital Sky Survey (SDSS; Szkody et al. 2002, 2003, 2004, 2005, 2006), but as Gänsicke (2005) points out, the colors of the CVs turned up by SDSS range down to the cutoff used to trigger spectroscopic followup, so more presumably remain undiscovered. Patterson (1998) gives a judicious estimate of the space density as 10^{-5} pc^{-3} .

¹Based on observations obtained at the MDM Observatory, operated by Dartmouth College, Columbia University, Ohio State University, and the University of Michigan.

The space density of CVs is predicted to be an order of magnitude higher by the evolution scenario of cataclysmic binaries (Shara et al. 1986). Unravelling the long-term evolution of CVs is largely dependent on an accurate measure of their space density.

Essentially all CVs are X-ray sources at some level, so a great many CVs have been discovered as the optical counterparts of X-ray sources. Because the X-ray surveys have well-understood completeness limits, these surveys have the potential to put CV space density estimates on a firmer footing (though very weak X-ray emitters will still be missed). Schwöpe et al. (2002) presented new optical identifications of CVs in the Rosat Bright Source (RBS) list. For many of their sources they estimated the absolute magnitude M_V using an empirical relationship between M_V and the emission equivalent width of $H\beta$ (Patterson 1984; his eqn. 16). One source – RBS 0490 – had a very large $EW(H\beta) = 230 \text{ \AA}$, for which the Patterson relation predicts a quite faint $M_V = 13.4$. This, combined with an estimated $m_V = 16$, gave a distance of only 33 pc. RBS 0490 was one of two objects for which Schwöpe et al. (2002) estimated $d \sim 30$ pc, the other being RBS 1955. Applying the $V/V_{\max}$ method to their data set, Schwöpe et al. (2002) estimated a relatively high CV space density, of order $3 \times 10^{-5} \text{ pc}^{-3}$. However, the inference of this high space density rested almost entirely on these two objects.

Largely because of the influence these two objects exert on the the space density estimate, we undertook to characterize them and estimate their distances. A study of RBS 1955 will be published elsewhere; here we concentrate on RBS 0490, which lies in Eridanus at $\alpha = 3^h54^m10^s.3, \delta = -16^\circ52'50''$ (ICRS).

2. Observations

All our data are from the 2.4m Hiltner reflector at MDM Observatory on Kitt Peak, Arizona. Table 1 lists the observations.

For spectroscopy, we used the modular spectrograph and a 600 line mm^{-1} grating, giving $2.0 \text{ \AA pixel}^{-1}$ and about $3.5 \text{ \AA}$ resolution from 4300 to $7500 \text{ \AA}$, with severe vignetting toward the ends of this range. The detector was a 2048^2 SITe CCD with $24 \text{ }\mu\text{m}$ pixels. For the most part, compari-

son lamps were taken frequently to track telescope flexure. In more recent observing runs, we have abandoned that procedure in favor of using night-sky lines to track the zero point of the wavelength solution. This technique saves telescope time and greatly simplifies operations, and extensive cross-checks show essentially no loss of accuracy. We observed flux standard stars when the weather was clear. The data reduction and analysis procedures were for the most part similar to those described in Thorstensen et al. (2004). However, to extract one-dimensional spectra from the two-dimensional images, we used a new, original implementation of the algorithm described by Horne (1986) instead of the IRAF *apsun* task, the main advantage being better bad-pixel rejection.

We used the same SITe CCD for direct imaging as for the spectroscopy. The procedures were identical to those described by Thorstensen (2003); briefly, for the parallax measurement, we took sets of around a dozen short (~ 100 s) *I*-band exposures on each visit to the source, with *V*-band exposures included when conditions were photometric. We also took two sets of *UBVI* exposures in 2002 October. Reducing these using Landolt (1992) standard star observations and mean extinction coefficients yielded the standard magnitudes given in Table 2. The coordinates in Table 2 are from a fit to numerous USNO A2.0 (Monet et al. 1996) stars, and are estimated accurate to $\sim 0''.3$.

In addition to the *UBVI* images, we have four *VI* image pairs taken on two other visits to the source; these show RBS 0490 with $17.16 < V < 17.49$, and $0.25 < V - I < 0.43$. The mean *V* magnitude overall was near 17.4.

2.1. Photometric Variability

The filter photometry taken 2002 October showed RBS 490 varying significantly between adjacent sets of exposures (Table 2). Because the variability of this star appears not to have been studied, we derived differential time series photometry using our *I*-band parallax images. We shifted the differential magnitudes to approximately match the standard magnitudes, and found RBS 490 varying, apparently irregularly, in the range $16.7 < I < 17.6$. Field stars of comparable brightness generally appeared constant to ~ 0.03 mag. The observations were grouped in

short bursts over several years, and hence were not well-suited for period finding, but do show that RBS 490 varies significantly.

2.2. The Spectrum.

Fig. 1 shows the mean spectrum, which resembles that given by Schwöpe et al. (2002). The emission lines, which are listed in Table 3, are extraordinarily strong – the equivalent width of $H\alpha$ is more than 400 Å! $HeII\ \lambda 4686$ does appear, but with relatively modest strength. The spectrum resembles those shown by Schwöpe et al. (2002), but the lines all appear single-peaked, whereas they describe them as double-peaked.

2.3. Radial Velocities

To measure velocities we used convolution algorithms developed by Schneider & Young (1980) and Shafter (1983). $H\alpha$, $H\beta$, $HeI\ \lambda 5876$, and $HeI\ \lambda 6678$ all gave useful results. We experimented with different choices of convolution function (derivative of a Gaussian, or two antisymmetric Gaussian) and a large range of convolution-function width. For the most part the choices of line, function, and width gave broadly similar results. The most notable exception to this was the strongest line, $H\alpha$, which sometimes did not give any plausible signal in its radial velocities. For a fiducial velocity time series, we adopt an average of velocities of $H\beta$ and the two strongest HeI lines, measured by convolving with the derivative of a Gaussian, optimized for a 12 Å (full width half maximum) line width. Table 4 lists these radial velocities.

We searched for periodicities in the radial velocity time series by fitting general-purpose sinusoids at a range of trial frequencies, and then plotting $1/\chi^2$, where χ^2 is the average of the squared fit residuals normalized to their estimated counting-statistics uncertainties. For the stronger lines, counting statistics gave unrealistically small errors, so χ^2 tended to be much greater than unity.

Fig. 2 shows the resulting periodogram. There is a remarkable lack of significant periodicities across the range of ‘normal’ CV periods, i.e., at frequencies below ~ 20 cycle d^{-1} . However, there is a strong suggestion of a frequency near 30 cycle d^{-1} , that is, near $P = 46$ min. Although our data are fairly extensive, and cover a wide range of hour angle (see Table 1), the data are noisy

enough that the daily cycle count is ambiguous. Nonetheless, if we arbitrarily choose one of the possible frequencies and fold the velocities on that frequency, we obtain the result shown in Fig. 3. While the scatter is large, the modulation does appear significant. Table 5 gives the parameters of best-fit sinusoids at the strongest 30 cycle d^{-1} frequencies.

Some combinations of lines and measuring methods showed weak evidence of lower-frequency modulations more consistent with expectations for CV orbital periods, but these marginal detections were not consistent from line to line or method to method, so we have no confidence that any of them are real. The ~ 30 cycle d^{-1} frequency, by contrast, appeared consistently in all the lines separately, save for $H\alpha$ with some choices of measurement parameters.

We do not claim to have definitively established the existence of this modulation, but we think it is probably real. It is difficult to estimate a realistic false-alarm probability without a realistic model of the underlying noise process; in a limited sample, ‘red’ noise, for example, spurious low-frequency periodicities can appear significant. However, higher frequencies are less susceptible to this problem, because they must maintain coherence over many cycles to create an apparent signal in the periodogram.

2.4. Parallax

The astrometric solution includes 100 I-band images taken on eight different observing runs between 2002 October and 2006 January. Table 6 gives the information for the 44 measured stars, of which 16 were used to establish the reference frame.

Table 7 lists the astrometric parameters. The parallax is barely detected, indicating a large distance, but the proper motion is very large, indicating a small distance. The Bayesian distance-estimation formalism explained in Thorstensen (2003) uses all this information. This formalism combines the measured proper motion with an *assumed* velocity distribution in order to truncate the divergence in probability at small parallaxes resulting from the Lutz-Kelker bias²; this diver-

²An absolute-magnitude constraint can also be applied, but this was not used here, because we have no prior knowledge

gence becomes severe when the relative error in the parallax is large, as it is here. The space velocity distribution assumed in the Bayesian analysis was the same as that used in Thorstensen (2003); it assumes that the bulk of CVs have a velocity distribution similar to disk stars, but that there is also a high-velocity tail. Because the model assumes that most CVs have rather low space velocities, a distance estimate based on the model and proper motion alone yields a 50th-percentile distance of only 58 pc; however, the small parallax firmly excludes any distance this short. At the distance indicated by the parallax, RBS 490 evidently lies in or near the high-velocity tail of the assumed distribution. The combination of all the evidence in the Bayesian formalism yields a nominal distance of $285(+120, -105)$ pc (68 per cent confidence). A lower limit to the distance can be established by ignoring the Lutz-Kelker correction and the Bayesian analysis, and taking $\pi_{\text{abs}} < \pi_{\text{abs}}(\text{observed}) + 2\sigma_{\pi} = 7.5$ mas; this exercise gives $d > 133$ pc. The Bayesian analysis gives a one-sigma upper limit distance of 405 pc. This would imply a transverse velocity near 200 km s^{-1} , which seems unlikely, since CVs generally have modest mean radial velocities γ (van Paradijs, Augusteijn, & Stehle 1996). Also, in RBS 0490, the γ velocity is $\sim 16 \text{ km s}^{-1}$; in an isotropic velocity distribution, a 200 km s^{-1} space velocity will result in $|\gamma| < 16 \text{ km s}^{-1}$ with probability ~ 0.08 . Thus $d < 405$ pc appears to be a reasonable upper limit. Because the Bayesian 50th percentile estimate of $d = 285$ pc depends strongly on the uncertain high-velocity tail of the CV distribution, we use 200 pc as a nominal distance, but we emphasize that this is very uncertain.

3. Discussion

3.1. Orbital Period

Ordinarily in CV studies, an emission-line radial velocity period can be confidently identified with the orbital period. Our apparent ~ 46 min period challenges that assumption.

CVs with ‘normal’ hydrogen-burning secondary stars are thought to evolve to shorter periods as they lose angular momentum, pass through a period minimum near $P_{\text{orb}} \sim 70$ min, and then

evolve toward longer periods (Patterson 1998). This is called the ‘period bounce’ (although it does not take place as abruptly as that term suggests). Most of the relatively small number of known CVs with periods below the ‘bounce’ show no hydrogen lines and appear to have helium secondaries; these are the AM CVn stars. Of the small handful of CVs with periods below the period minimum that do not obviously have helium secondaries, at least one – EI Psc — has an anomalously warm secondary that is evidently somewhat evolved (Thorstensen et al. 2002a), and the helium lines in both EI Psc and the 59-minute binary V485 Cen (Augusteijn et al. 1996) appear at least a little strong when compared with the ordinary hydrogen-accretors at longer periods. In RBS 0490 the emission lines are remarkably strong, but the He lines do not appear to be enhanced, so an orbital period as short as 46 minutes seems implausible.

In recent years, several objects have been discovered that show radial velocity periods which are clearly not the same as P_{orb} . The emission velocities in HS 2331+3905 show a remarkably convincing quasi-period near 3.5 hr (Araujo-Betancor et al. 2005), but the periodicity does not retain coherence over longer timescales, and weak eclipses indicate a stable a P_{orb} near 81 min. In DW Cnc, the emission-line radial velocities reveal stable periodicities at 86.1 and 38.6 min, with comparable amplitude; it appears that these periodicities represent respectively the orbit and the white-dwarf spin (Patterson et al. 2004; Rodríguez-Gil et al. 2004). These examples add weight to the argument that the ~ 46 min period in RBS 0490 is not orbital.

But, if the 46-min period is *not* P_{orb} , then the absence of any radial-velocity modulation at a more convincing orbital frequency is somewhat surprising; while CV emission lines often do not trace the motion of the white dwarf accurately, they usually do show some modulation at P_{orb} . The full-width at half maximum of the $\text{H}\alpha$ emission line is 1000 km s^{-1} , so it is unlikely that the system is nearly face-on. It is possible that the secondary’s mass is very small. The spectrum shows no trace of the secondary, but given the still-unknown P_{orb} and rather uncertain distance, the non-detection is not readily interpreted physically.

Fortunately, rather than crying vaguely for

of this system’s luminosity.

‘more studies’, we can suggest a very specific approach to unscramble this object: extensive time-series photometry. There is a good chance that this would reveal other periodicities that would point the way toward understanding this object.

3.2. Distance and Demographics

Our mean $V = 17.4$ implies an absolute magnitude $M_V = 10.9 - 5 \log_{10}(d/200 \text{ pc})$, or $M_V < 11.8$ using our lower-limit distance of 133 pc. As noted earlier, the M_V -EW(H β) relation from Patterson (1984) predicts $M_V = 13.4$ from the very strong emission. However, the relationship shows considerable scatter, and the calibration points used by Patterson (1984) do not extend to objects with lines this strong. One must clearly be cautious in using this broad-brush correlation to estimate the luminosities of individual objects; also, the lines in this object really do appear extraordinarily strong for the luminosity. The very short distance estimated by Schwöpe et al. (2002) was also based on a value of $V \sim 16$, estimated from spectrophotometry; this is over 1 magnitude brighter than we have seen it in our four photometric visits. In any case, the short distance is excluded by the small parallax. Also, an MDM Observatory parallax study of RBS 1955 (in preparation) gives a distance of ~ 160 pc for that source, over five times the Schwöpe et al. (2002) estimate. Thus it appears that the high space density derived in that paper should be disregarded.

The appropriate CV subclassification for RBS 0490 is unclear. The relatively short period in the emission lines suggests a white dwarf spin period in a magnetic system, but HeII $\lambda 4686$ is rather weak, while in most magnetic CVs it has considerable strength. On the other hand, HeII $\lambda 4686$ is not particularly strong in DW Cnc or HS 2331+3905, either, so this is not conclusive. The absolute magnitude is comparable to those of dwarf novae in quiescence, and much less bright than novalike variables. Using our upper-limit distance of 405 pc, we find $M_V > 9.4$, which is still well short of novalike brightness.

Acknowledgments. We gratefully acknowledge support through NSF grants AST 9987334 and AST 0307413, and thank Bill Fenton for taking some of the spectroscopic data.

REFERENCES

- Araujo-Betancor, S., et al. 2005, A&A, 430, 629
- Augusteijn, T., van der Hooft, F., de Jong, J. A., & van Paradijs, J. 1996, A&A, 311, 889
- Gänsicke, B. T. 2005, ASP Conf. Ser. 330: The Astrophysics of Cataclysmic Variables and Related Objects , 330, 3 764
- Horne, K. 1986, PASP, 98, 609
- Landolt, A. U. 1992, AJ, 104, 340
- Monet, D., et al. 1996, USNO-SA2.0, (U. S. Naval Observatory, Washington, DC)
- Patterson, J. 1984, ApJS, 54, 443
- Patterson, J. 1998, PASP, 110, 1132
- Patterson, J., et al. 2004, PASP, 116, 516
- Pojmański, G. 2001, ASP Conf. Ser. 246: IAU Colloq. 183: Small Telescope Astronomy on Global Scales, 246, 53
- Rodríguez-Gil, P., Gänsicke, B. T., Araujo-Betancor, S., & Casares, J. 2004, MNRAS, 349, 367
- Schwope, A. D., Brunner, H., Buckley, D., Greiner, J., Heyden, K. v. d., Neizvestny, S., Potter, S., & Schwarz, R. 2002, A&A, 396, 895
- Schneider, D. and Young, P. 1980, ApJ, 238, 946
- Shafter, A. W. 1983, ApJ, 267, 222
- Shara, M. M., Livio, M., Moffat, A. F. J., & Orio, M. 1986, ApJ, 311, 163
- Szkody, P., et al. 2002, AJ, 123, 430
- Szkody, P., et al. 2003, AJ, 126, 1499
- Szkody, P., et al. 2004, AJ, 128, 1882
- Szkody, P., et al. 2005, AJ, 129, 2386
- Szkody, P., et al. 2006, AJ, 131, 973
- Thorstensen, J. R. 2003, AJ, 126, 3017
- Thorstensen, J. R., Fenton, W. H., Patterson, J. O., Kemp, J., Krajci, T., & Baraffe, I. 2002, ApJ, 567, L49
- Thorstensen, J. R., Fenton, W. H., & Taylor, C. J. 2004, PASP, 116, 300
- van Paradijs, J., Augusteijn, T., & Stehle, R. 1996, A&A, 312, 93
- Warner, B. 1995, Cataclysmic Variables (Cambridge University Press)

TABLE 1
JOURNAL OF OBSERVATIONS

Date [UT]	N	HA (start) [hh:mm]	HA (end) [hh:mm]	p_X^a [arcsec]	p_Y [arcsec]
Parallax Observations:					
2002 Oct 24	1	−2:08	−2:08	0.49	−0.44
2003 Jan 29	15	+0:29	+1:28	−0.89	−0.37
2004 Jan 10	8	−0:13	+0:15	−0.72	−0.51
2004 Feb 29	14	+1:33	+2:04	−0.95	−0.06
2004 Nov 12	14	−0:53	+1:12	0.17	−0.56
2005 Jan 30	7	+0:18	+0:39	−0.90	−0.36
2005 Sep 13	24	−0:19	+0:28	0.92	−0.06
2006 Jan 15	17	−0:08	+1:47	−0.78	−0.48
Spectroscopy:					
2002 Oct 26	10	−2:47	+3:26		
2002 Oct 29	33	−2:16	+3:22		
2002 Oct 30	25	−2:24	+2:56		
2002 Oct 31	34	−3:01	+3:15		
2002 Dec 12	3	−0:59	−0:48		
<i>UVBI</i> photometry (two image sets):					
2002 Oct 25	8	−0:15	+0:14		

^aParallax factor in X (right ascension); p_Y is the parallax factor in declination.

TABLE 2
FILTER PHOTOMETRY

α^a	δ^a	$U - B$	$B - V$	V	$V - I$
Field stars (averages of two observations)					
3:53:56.04	−16:52:08.0	0.26 ± 0.02	0.48 ± 0.01	16.69 ± 0.01	0.78 ± 0.01
3:54:01.07	−16:54:53.4	0.31 ± 0.01	0.73 ± 0.01	16.16 ± 0.01	0.78 ± 0.01
3:54:01.62	−16:52:22.9	-0.27 ± 0.06	0.50 ± 0.03	18.72 ± 0.02	0.70 ± 0.02
3:54:02.88	−16:51:16.3	-0.21 ± 0.05	0.44 ± 0.02	18.55 ± 0.02	0.72 ± 0.02
3:54:04.02	−16:49:48.0	...	1.74 ± 0.07	18.80 ± 0.02	3.16 ± 0.02
3:54:05.06	−16:49:14.7	...	1.01 ± 0.08	19.59 ± 0.04	0.96 ± 0.05
3:54:05.50	−16:55:26.3	...	1.36 ± 0.04	18.45 ± 0.01	2.20 ± 0.02
3:54:06.05	−16:54:29.4	...	1.54 ± 0.13	19.66 ± 0.04	1.99 ± 0.05
3:54:06.35	−16:53:17.1	0.07 ± 0.01	0.55 ± 0.01	14.77 ± 0.01	0.64 ± 0.01
3:54:06.49	−16:50:25.9	...	1.48 ± 0.06	18.89 ± 0.02	2.38 ± 0.02
3:54:06.97	−16:50:44.9	0.02 ± 0.01	0.62 ± 0.01	15.99 ± 0.01	0.71 ± 0.01
3:54:06.98	−16:50:39.9	-0.00 ± 0.01	0.60 ± 0.01	15.86 ± 0.01	0.69 ± 0.01
3:54:08.80	−16:54:45.9	0.55 ± 0.01	0.82 ± 0.01	15.70 ± 0.01	0.88 ± 0.01
3:54:10.15	−16:53:05.4	...	0.63 ± 0.02	18.15 ± 0.01	0.75 ± 0.02
3:54:12.45	−16:51:31.3	-0.91 ± 0.06	-0.07 ± 0.05	19.58 ± 0.04	0.85 ± 0.05
3:54:13.59	−16:52:27.3	0.48 ± 0.09	0.96 ± 0.02	18.03 ± 0.01	1.01 ± 0.01
3:54:15.22	−16:49:04.0	1.04 ± 0.17	1.52 ± 0.03	17.80 ± 0.01	1.90 ± 0.01
3:54:18.50	−16:51:25.2	...	1.45 ± 0.03	17.99 ± 0.01	1.68 ± 0.01
3:54:24.08	−16:55:24.7	...	1.20 ± 0.03	18.49 ± 0.02	1.47 ± 0.02
Variable star (two observations separately)					
3:54:10.30	−16:52:49.8	-1.26 ± 0.01	-0.24 ± 0.01	17.58 ± 0.01	0.55 ± 0.01
...	...	-1.32 ± 0.01	-0.17 ± 0.01	17.71 ± 0.01	0.77 ± 0.01

^aCoordinates referred to the ICRS, and are from a fit to 24 USNO A2.0 stars, with a scatter of 0.6 arcsec. Units are hours, minutes, and seconds for the right ascension and degrees, minutes, and seconds for the declination.

TABLE 3
EMISSION FEATURES

Feature	E.W. ^a (Å)	Flux ^b	FWHM ^c (Å)
HeI λ 4471	27	97	13
HeII λ 4686	12	34	15
HeI λ 4713	5	13	20
H β	250	720	17
HeI λ 4921	15	40	18
HeI λ 5015	11	26	15
HeI λ 5876	95	160	18
H α	430	750	22
HeI λ 6678	48	78	20
HeI λ 7067	33	48	22
HeI λ 7281	9	13	22

^aEmission equivalent widths are counted as positive.

^bUnits are 10^{-16} erg cm $^{-2}$ s $^{-1}$. Absolute line fluxes are uncertain by about 30 per cent, but relative fluxes of strong lines should be accurate to ~ 10 per cent.

^cFrom Gaussian fits.

TABLE 4
RADIAL VELOCITIES

Time ^a	v_{em}^{b} (km s ⁻¹)	Time ^a	v_{em}^{b} (km s ⁻¹)	Time ^a	v_{em}^{b} (km s ⁻¹)
573.7652	-17	576.9505	100	578.7516	-35
573.7714	-8	576.9845	-18	578.7564	-90
573.7775	3	576.9892	14	578.7611	59
573.7836	19	576.9940	-51	578.7659	119
573.9604	78	576.9987	-63	578.7706	31
573.9665	28	577.0035	-64	578.7784	-22
574.0022	-14	577.0082	4	578.7831	65
574.0079	-42	577.0130	-38	578.7879	47
574.0174	20	577.7703	-40	578.7926	107
574.0238	37	577.7751	-28	578.7974	47
576.7785	26	577.7798	49	578.8022	-17
576.7830	75	577.7846	99	578.9030	-18
576.7881	-14	577.7894	24	578.9077	-55
576.7932	-68	577.7941	3	578.9125	-64
576.7979	-84	577.7989	-35	578.9172	51
576.8082	30	577.9058	-70	578.9220	137
576.8149	18	577.9106	-1	578.9267	102
576.8196	-74	577.9155	-6	578.9315	26
576.8244	2	577.9203	96	578.9383	-20
576.8291	-37	577.9250	84	578.9431	-11
576.8339	30	577.9298	-5	578.9478	44
576.8823	81	577.9345	-20	578.9526	14
576.8867	75	577.9397	12	578.9574	2
576.8910	5	577.9467	55	578.9621	25
576.8954	-60	577.9515	133	578.9669	6
576.9003	7	577.9562	68	578.9742	-80
576.9056	75	577.9610	-34	578.9789	-44
576.9104	94	577.9657	-71	578.9837	64
576.9151	8	577.9705	-40	578.9884	58
576.9220	-62	577.9753	30	578.9932	96
576.9267	72	577.9822	-3	578.9980	31
576.9315	78	577.9870	-45	579.0027	-1
576.9362	52	577.9918	28	620.7112	26
576.9410	83	578.7421	54	620.7153	83
576.9458	70	578.7469	31	620.7193	143

^aHeliocentric Julian date of mid-exposure, minus 2 452 000.

^bAverage velocity of H β , HeI λ 5876, and HeI λ 6678.

TABLE 5
FIT TO THE RADIAL VELOCITIES

T_0^{a}	P (d)	K (km s ⁻¹)	γ (km s ⁻¹)	N	σ^{b} (km s ⁻¹)
577.7755(16)	0.03356(3)	46(13)	16(9)	102	42
577.7793(16)	0.03247(4)	44(13)	17(9)	102	43
577.771(2)	0.03472(5)	39(14)	16(10)	102	46

NOTE.—Parameters of least-squares sinusoid fits to the radial velocities, of the form $v(t) = \gamma + K \sin(2\pi(t - T_0)/P)$.

^aHeliocentric Julian Date minus 2452000.

^bRoot-mean-square residual of the fit.

TABLE 6
ASTROMETRIC RESULTS

α^a [hh:mm:ss]	δ [dd:mm:ss]	W^b	σ^c [mas]	V	$V - I$	π_{rel} [mas]	μ_X^d	μ_Y [mas yr $^{-1}$]	σ_μ
3:53:55.81	-16:52:54.4	1	6	17.86	2.15	2.4 ± 0.8	-3.3	-4.5	0.6
3:53:56.04	-16:52:08.4	1	6	16.69	0.80	-3.0 ± 0.7	-2.3	4.2	0.6
3:53:56.91	-16:55:21.8	0	32	21.52	2.58	-10.5 ± 3.8	11.4	-18.5	3.2
3:53:59.05	-16:54:03.0	0	63	20.55	0.71	-0.2 ± 7.4	4.4	-15.5	6.3
3:54:01.08	-16:54:53.9	1	6	16.17	0.80	-0.7 ± 0.7	2.6	0.6	0.6
3:54:01.48	-16:53:21.0	0	20	21.01	2.72	-2.2 ± 2.4	11.1	-0.9	2.0
3:54:01.62	-16:52:23.4	0	17	18.73	0.68	-6.8 ± 2.0	-1.5	-2.3	1.7
3:54:02.30	-16:54:08.2	0	52	20.98	1.40	0.4 ± 6.1	-1.0	-4.9	5.2
3:54:03.20	-16:54:22.3	0	145	19.80	1.12	22.1 ± 17.0	-16.7	0.3	14.5
3:54:02.88	-16:51:16.8	0	12	18.51	0.68	-2.4 ± 1.5	-7.3	4.1	1.2
3:54:03.02	-16:51:47.3	0	47	22.01	2.40	3.6 ± 5.5	-10.0	-12.8	4.7
3:54:04.37	-16:54:32.0	0	48	22.16	2.64	-4.5 ± 5.7	21.4	2.1	4.8
3:54:04.09	-16:50:49.5	0	10	18.44	1.43	-3.7 ± 1.2	-14.4	10.0	1.0
3:54:04.02	-16:49:48.4	1	7	18.71	3.04	1.2 ± 0.8	24.6	-0.8	0.7
3:54:05.50	-16:55:26.8	1	6	18.43	2.17	0.5 ± 0.7	4.5	0.2	0.6
3:54:05.06	-16:49:15.0	0	28	19.65	0.94	-4.6 ± 3.2	-43.5	21.4	2.8
3:54:06.05	-16:54:29.9	0	15	19.64	1.90	-3.7 ± 1.8	15.9	-4.9	1.5
3:54:06.50	-16:54:29.7	0	30	19.63	0.75	-7.9 ± 3.5	8.7	-2.2	3.0
3:54:06.36	-16:53:17.5	1	8	14.77	0.64	-1.0 ± 0.9	12.2	-3.1	0.8
3:54:06.49	-16:50:26.4	1	7	18.88	2.33	-2.0 ± 0.8	-5.3	-9.8	0.7
3:54:06.98	-16:50:45.4	1	5	15.98	0.72	-0.8 ± 0.6	-10.6	9.1	0.5
3:54:06.98	-16:50:40.3	1	5	15.85	0.70	-0.6 ± 0.6	-11.6	8.9	0.5
3:54:07.35	-16:51:03.8	1	10	20.52	2.91	2.0 ± 1.1	-14.7	-9.1	0.9
3:54:08.52	-16:54:24.8	0	15	20.36	2.43	0.4 ± 1.8	1.0	-4.0	1.5
3:54:08.81	-16:54:46.4	1	8	15.70	0.88	-1.0 ± 0.9	2.1	2.1	0.8
3:54:08.43	-16:51:37.0	0	26	20.12	1.33	1.1 ± 3.1	-4.1	2.8	2.6
3:54:08.79	-16:51:19.7	0	29	19.67	0.72	1.5 ± 3.4	-3.0	-0.8	2.9
3:54:09.50	-16:52:24.0	0	28	21.14	2.32	1.9 ± 3.2	3.5	5.4	2.8
3:54:10.00	-16:54:25.8	0	19	20.81	2.48	-0.8 ± 2.2	4.0	6.5	1.9
3:54:10.16	-16:53:05.9	1	9	18.18	0.76	0.9 ± 1.1	5.1	1.1	0.9
3:54:10.31	-16:52:50.3	0	9	17.41	0.39	3.2 ± 1.0	-2.5	-102.3	0.9
3:54:11.37	-16:55:34.4	1	5	20.75	2.89	0.2 ± 0.6	-12.6	-1.6	0.5
3:54:11.42	-16:53:23.4	0	32	21.69	2.45	-0.5 ± 3.7	17.0	-8.6	3.2
3:54:12.46	-16:51:31.8	0	24	19.65	0.81	2.4 ± 2.8	-4.1	-0.4	2.4
3:54:13.60	-16:52:27.7	1	9	18.07	1.03	-1.4 ± 1.0	2.4	1.8	0.9
3:54:15.62	-16:55:32.1	0	24	19.88	1.35	-4.5 ± 2.8	-7.8	23.7	2.4
3:54:15.22	-16:49:04.4	0	17	17.81	1.90	0.2 ± 2.0	-25.2	11.9	1.7
3:54:18.51	-16:51:25.6	1	9	18.00	1.69	1.2 ± 1.1	7.9	-2.1	0.9
3:54:19.99	-16:54:56.9	0	43	22.52	3.23	1.3 ± 5.1	6.2	6.6	4.3
3:54:24.10	-16:55:25.2	0	40	18.45	1.43	-0.1 ± 4.7	0.5	30.5	4.0
3:54:24.07	-16:50:52.5	0	40	22.36	2.85	-0.4 ± 4.7	16.8	-3.8	4.0
3:54:24.32	-16:50:13.4	0	47	21.71	2.53	-5.5 ± 5.6	4.9	-5.4	4.7
3:54:26.17	-16:50:45.2	0	29	20.56	1.99	-4.8 ± 3.4	9.5	-9.7	2.9
3:54:25.90	-16:49:10.3	1	3	16.74	0.91	-0.1 ± 0.4	-1.9	0.3	0.3

^aReferred to the ICRS ($\sim$ J2000).

^bWeight, 1 or 0 depending on whether the star is used as reference.

^cRMS deviation from the relative positions predicted by the best-fit parallax and proper motion.

^dProper motions are relative to the set of reference stars used, and are not on an inertial system.

NOTE.—Astrometric parameters for all the stars measured in the field. The program star is marked with an asterisk.




Fig. 1.— Mean spectrum of RBS490.




Fig. 2.— Periodogram of the fiducial radial velocity time series.




Fig. 3.— Emission-line radial velocities, folded on one of the 30 cycle d^{-1} alias periods, together with the best-fitting sinusoid. The data and fit are repeated for a second cycle to preserve continuity. The open squares show velocities from 2002 December, and the rest of the symbols are from 2002 October. The fold is based on an arbitrary choice of cycle counts between these two epochs.